



TREATISE ONLINE

Number 109

Part V, Second Revision, Chapter 2:
Class Enteropneusta: Introduction, Morphology,
Life Habits, Systematic Descriptions, and
Future Research

Christopher B. Cameron

2018

KU PALEONTOLOGICAL
INSTITUTE

The University of Kansas

Lawrence, Kansas, USA
ISSN 2153-4012
paleo.ku.edu/treatiseonline

PART V, SECOND REVISION, CHAPTER 2: CLASS ENTEROPNEUSTA: INTRODUCTION, MORPHOLOGY, LIFE HABITS, SYSTEMATIC DESCRIPTIONS, AND FUTURE RESEARCH

CHRISTOPHER B. CAMERON

[Département de sciences biologiques, Université de Montréal, Montréal QC, H2V 2S9, Canada, c.cameron@umontreal.ca]

Class ENTEROPNEUSTA Gegenbaur, 1870

[*nom. correct.* HAECKEL, 1879, p. 469 *pro* Enteropneusti GEGENBAUR, 1870, p. 158]

Free living, solitary, worms ranging from lengths of less than a millimeter to 1.5 meters; entirely marine; body tripartite, with proboscis, collar, and trunk; proboscis coelom contains heart-kidney-stomochord complex; preoral ciliary organ posterior-ventral; collagenous Y-shaped nuchal skeleton extends from proboscis through neck before bifurcating into paired horns in collar; paired dorsal periaemal coeloms associated with collar dorsal blood vessel; anterior trunk pharynx perforated with paired gill slits that connect via atria to external gill pores; mesocoel ducts connect collar coeloms to first pair of gill pores atria; larvae possess locomotory ciliated band (telotroch), and when tornaria present, apical plate retractor muscle. *Cambrian (Miaolingian, Wuliuan)–Holocene* (extant): worldwide.

The class Enteropneusta is differentiated among four living families: the Harrimaniidae, Spengelidae, Ptychoderidae, and Torquaratoridae, based in large part on the structure of the heart-kidney coelomic complex, gills, gonads, liver sacs, and diverticula of the coelomic compartments. The modern families are a crown group clade with respect to the Cambrian fossil stem group, which includes *Oesia disjuncta* and *Spartobranchus tenuis* (see p. 12).

MORPHOLOGY

The acorn worm body is arranged into an anterior proboscis, a collar, and a posterior trunk (Fig. 1). Body length can vary from less than a millimeter (WORSAAE & others, 2012) to 1.5 meters (SPENGLER, 1893). The proboscis is muscular and its epidermis replete with sensory, ciliated, and glandular cells (BENITO & PARDOS, 1997). Acorn worms deposit-feed by trapping sediment in mucus and transporting it to the mouth with cilia. A pre-oral ciliary organ on the posterior proboscis (BRAMBEL & COLE, 1939) (Fig. 2.1) directs the food-laden mucous thread into the mouth (GONZALEZ & CAMERON, 2009). The proboscis coelom contains a turgid stomochord (Fig. 1, Fig. 2.2a), an anterior projection of the dorsal collar buccal cavity that consists of vacuolated cells arranged around a central ciliated cavity and encapsulated in a collagen sheath (WELSCH & STORCH, 1970). The result is a rigid structure against which the pericardium (heart) compresses the intervening blood sinus so that urine can be forced across the glomerulus (kidney) into the proboscis coelom (BALSER & RUPPERT, 1990). The stomochord is not a homolog of the chordate notochord (SATOH & others, 2014). The right protocoeleum is small or absent, and the left coelom is dominant and extends a ciliated duct, which is lined with nephridia (mesothelial monociliated podocytes and myocytes) to the exterior via




FIG. 1. Illustration of a generalized morphology of an enteropneust. Anus (*an*), atrium (*at*), dorsal nerve cord (*dc*), dorsal blood vessel (*dv*), gonad (*gd*), filtration glomerulus (*gl*), gill pore (*gp*), gill slit (*gs*), heart sinus (*hs*), liver sac (*ls*), mesocoel (*mc*), paired mesocoel ducts (*md*), mouth (*mo*), metacoels (*mt*), Y-shaped nuchal skeleton (*ns*), peribuccal coeloms (*pb*), protocoele (*pc*), pericardium (*pe*), ventral post-anal tail (*pt*), preoral ciliary organ (*poco*), urinary pore (*pp*), periahaemal coeloms (*pr*), stomochord (*st*), and ventral blood vessel (*vv*) (adapted from Cameron 2005, fig. 1).

a left dorsal lateral urinary, proboscis pore (RUPPERT & BALSER, 1986) (Fig. 1). There is extensive inter- and intraspecific variation in the form of this proboscis heart-kidney-coelom complex (DELAND & others, 2010). The stomochord is supported ventrally by a collagenous Y-shaped nuchal skeleton (Fig. 1, Fig. 2.2a–b). The skeleton extends posteriorly through the neck and bifurcates into paired horns in the collar (SPENGLER, 1893; HORST, 1939; EZHOVA & MALAKHOV, 2009).

The collar lip forms the mouth anteriorly, and its interior is the buccal cavity (Fig. 1). In some species, the anterior lip is muscular and flexible, adept at guiding sediment laden strings of mucus from the proboscis into the mouth or rejecting them over the lip and onto the collar. The collar epidermis is usually differentiated into histologically distinct zones of transverse bands of numerous gland cells and dense bands of cilia (BENITO & PARDOS, 1997; HORST, 1939), with distinct mucin and glycoprotein secretory patterns (SIMAKOV & others, 2015). The acorn worm nervous system is an epidermal plexus with two longitudinal intraepidermal nerve cords in the trunk and a neurulated cord in the collar (BULLOCK, 1940; KNIGHT-JONES, 1953). Sensory cells, ganglion cells, interneuronal junctions, and motor innervation are arranged diffusely (BULLOCK, 1940; BENITO & PARDOS, 1997). The collar cord is intraepithelial even where the epithelium

containing the nerves becomes internalized, and most commonly has paired, parallel, and discontinuous lacuna that are lined with cilia. There is no central nervous system. The collar cord is a simple conduction pathway (CAMERON & MACKIE, 1996) and is aligned with the periahaemal muscles, suggesting a role in the rapid contraction of the proboscis (RUPPERT, 2005). This cord is unlike the hollow, neuroepithelium found in the rays of echinoderms, or the hollow, neural tube of chordates (BULLOCK, 1940; CAMERON & MACKIE, 1996; RUPPERT, 2005). Paired diverticula of the trunk coeloms, called periahaemal and peribuccal coeloms, project anteriorly into the collar (Fig. 1). Paired mesocoel ducts connect the collar coeloms to the first pair of gill pore atria in the anterior trunk (Fig. 1).

The acorn worm trunk bears middorsal and midventral longitudinal grooves that correspond to the location of longitudinal nerves and blood vessels (PARDOS & BENITO, 1990) (Fig. 1). The dorsal and ventral nerve cords are connected anteriorly by a circumpharyngeal ring (BULLOCK, 1940; KAUL-STREHLOW & others, 2015). The ventral cord is the fastest conducting cord due to its proximity to the large ventral longitudinal muscles (PICKENS, 1970; CAMERON & MACKIE, 1996). The trunk is differentiated into regions along its length that correspond to the specializations of the gut. The anterior

region bears paired dorsolateral series of gills, followed by an esophagus, and an intestine that terminates at an anus (Fig. 1). The gill slits, demarcated by a series of M-shaped collagenous gill bars, are connected to the outside pores by atrial cavities, akin to those of cephalochordates. The ciliated bars pump water through the pharynx, slits, and pores, removing food particles from suspension (CAMERON, 2002a; GONZALEZ & CAMERON, 2009). Support for the hypothesis that gills are a deuterostome plesiomorphy are shared patterns of genes expressed during gill and pharynx development and the organization of these genes in synteny in acorn worms and vertebrates (GILLIS, FRITZENWANKER, & LOWE, 2012; SIMAKOV & others, 2015). The food and mucus are then transported down the gill bars to the ventral gut, concentrated into a mucous food cord, and then transported posteriorly through the esophagus. Iodine¹²⁵ binding experiments suggests that an endostyle function may reside in the entire pharyngeal lining of Enteropneusta, and in *Schizocardium* SPENGLER, 1893, an epibranchial ridge is organized into zones of cells, reminiscent of the chordate endostyle (RUPPERT, CAMERON, & FRICK, 1999; RUPPERT, 2005; see also SATOH & others, 2014) (Fig. 2.3). The muscular esophagus functions to eliminate excess water from the food cord as it passes to the intestine. Anteriorly, the intestine is usually darkly pigmented. It is a simple tube except in *Schizocardium* and the family Ptychoderidae, in which it develops dorsally projecting rows of hepatic sacs, presumably to deal with a diet of heavy sediment. Gonads occur along the pharyngeal region, and in the family Ptychoderidae, they develop in external paired longitudinal ridges or wings. A few species have a curious pygochord structure in the posterior ventral trunk that WILLEY (1899) suggested may be a homolog to the chordate notochord. It is comprised of vacuolated cells positioned between the posterior ventral trunk mesenteries (WILLEY, 1899; SPENGLER, 1903; URIBE & LARRAIN, 1992; MIYAMOTO & SAITO, 2007) or intestine

(HORST, 1939). PUNNETT (1906) suggested that the structure may be a vestigial collateral intestine, which may instead function to support the fragile trunk as it projects from the burrow to eject fecal sediment. Distinct protuberances characterize the epidermis of the acorn worm trunk. These are formed by secretory and sclerocyte cells that occlude extracellular spaces, where microscopic calcium carbonate ossicles develop. In *Ptychodera* (Fig. 2.4a) and *Saccoglossus* (Fig. 2.4b), the ossicles form a polycrystalline aggregate of laminar layers, with perforations that are comparable to the echinoderm stereome (CAMERON & BISHOP, 2012). The functions of the pygochord, and the ossicles are unknown.

DEVELOPMENT AND REPRODUCTION

The sexes are separated except for *Yoda purpurata* PRIEDE & others, 2012 which is a hermaphrodite (see PRIEDE & others, 2012). Fertilization is external (BURDON-JONES 1951; LOWE & others 2004) although vivipary is known for *Xenopleura vivipara* GILCHRIST, 1925, and the deep-sea *Coleodesmium karaensis* OSBORN & others, 2013, which broods their embryos in sacs that connect to a gonadal wing by a narrow stock (OSBORN & others, 2013). In most species, eggs are released in a jelly coating that breaks down following fertilization (COLWIN & COLWIN, 1954). All subsequent development occurs in the seawater. Cleavage is holoblastic, radial, and more or less equal. Cell fates and axis specifications are in most respects similar to that of an echinoderm (HENRY, TAGAWA, & MARTINDALE, 2001). A coeloblastula is followed by gastrulation by invagination. The blastopore is located posteriorly but closes before a new anus and mouth are formed (BURDON-JONES, 1952; COLWIN & COLWIN, 1953). The fertilization envelope weakens and ruptures, releasing ciliated, swimming embryos by late gastrula. The organization of the coelomic sacs is comparable to echinoderms. The ambulacrarian coeloms are organized anterior to posterior as




FIG. 2. Acorn worm anatomy. 1, Pre-oral ciliary organ of *Protoglossus graveolens* GIRAY & KING, 1996; 2a, stomochord, nuchal skeleton, and gill bars of *P. graveolens*; 2b, illustrations of nuchal skeletons, in clockwise order, *Protoglossus koehleri* (CAULLERY & MESNIL, 1900), *Saccoglossus ruber* TATTERSALL, 1905, and *Balanoglossus hydrocephalus* HORST, 1940; 3, epibranchial ridge of *Schizocardium peruvianum* SPENGLER, 1893; 4a, ossicles of *Ptychodera flava* var. ESCHSCHOLTZ, 1825, Galapagos; 4b, ossicles of *Saccoglossus bromophenolus* KING, GIRAY, & KORNFIELD, 1994 (all, new).




FIG. 3. SEM photos of details. 1, gill region of *Harrimania planktophilus* CAMERON, 2002b; 2, gill region of *Ptychodera* sp.; 3, *Saccoglossus kowalevskii* AGASSIZ, 1873 at the three-gill pore stage, showing a typical harrimaniid juvenile post-anal tail; 4, posteriofrontal view of a tornaria larva of *Schizocardium* sp., showing the swimming telotrochal band (all, new).

paired protoceol, mesocoels, and metacoels (CRAWFORD & CHIA, 1978). They develop as independent pouches from the archenteron by enterocoely (KAUL-STREHLOW & STACH, 2013). Reports of schizocoely have not been substantiated (HADFIELD, 1975). GEMMILL (1914) provides a thorough comparative study of coelomogenesis in the asteroid (starfish), *Asterias rubens* LINNAEUS, 1758 and an enteropneust (summarized in CAMERON, 2005).

Harrimaniid worms (e.g., *Saccoglossus*) develop from a large, yolky egg to a ciliated, non-feeding larva with a telotrochal band and apical tuft that allows for a brief swimming and exploratory period (COLWIN & COLWIN, 1953). Following settlement, the juvenile worm has a ciliated, adhesive,

post-anal tail (Fig. 3.3) that allows for rapid forward and reverse locomotion, adhesion, and release (BURDON-JONES, 1952; BURDON-JONES 1956; CAMERON, 2002b; STACH & KAUL, 2011) that may be a homolog to the pterobranch stalk (BURDON-JONES, 1952; HYMAN, 1959; LESTER, 1985; CAMERON, 2005). Ptychoderid worms (e.g., *Ptychodera*) shed small transparent eggs. Development is rapid and indirect via a tornaria larva that resembles an echinoderm auricularia (MORGAN, 1891; GEMMILL, 1914; GISLÉN, 1930; TAGAWA & others, 1998; URATA & YAMAGUCHI, 2004) (Fig. 3.4). These diplu- rula larvae are delicate and transparent with a preoral ciliated feeding band (Fig. 3.4) that creates an upstream feeding current using monociliated cells (MORGAN, 1891;

STRATHMANN & BONAR, 1976), and a perioral ciliated band that manipulates and conveys food into the esophagus (LACALI & GILMOUR, 2001). The free-swimming, feeding tornaria are able to turn due to a band of multiciliated cells that comprise the locomotory telotroch (Fig. 3.4), an acorn worm innovation. The tornaria apical plate retractor muscle (and associated coughing behavior) (MORGAN, 1891) is another acorn worm larval apomorphy. Tornaria may persist in the plankton for months before metamorphosis and settlement to a suitable juvenile worm habitat. The larval body elongates differentiating the three body regions and organs. The larval apical tuft, nervous system, and feeding band are lost, followed by the telotroch band. The gills develop as out-pockets of the gut that open to the exterior by paired dorsolateral pores. Nothing is known of torquaratorid development, but the exceptionally large egg size (OSBORN & others, 2012; PRIEDE & others, 2012) hints that development is direct, via a loss of a feeding tornaria, parallel to the ontological loss of the larva in the line to modern harrimaniids.

Asexual reproduction by fragmentation is widespread among enteropneusts, including ptychoderids (GILCHRIST, 1923; PACKARD, 1968; MIYAMOTO & SAITO, 2010; HUMPHREYS & others, 2010), a spengelid (URATA, IWASAKI, & OHTSUKA, 2012), and the minute harrimaniid *Meioglossus psammophilus* WORSAAE & others, 2012, which lacks males and reproduces by paratomy. Regeneration is particularly well developed in the ptychoderids, a group especially susceptible to breakage due to their larger and fragile bodies. The posterior fragments are particularly good at regeneration, although some experimentally bisected animals can regenerate new individuals from both anterior and posterior fragments (WILLEY, 1899; DAWYDOFF, 1909, 1948; RAO, 1955; TWEDELL, 1961; PACKARD, 1968; PETERSEN & DITADI, 1971; RYCHEL & SWALLA, 2009; HUMPHREYS & others, 2010; MIYAMOTO & SAITO, 2010). The cells

that participate in blastema formation and morphogenesis during regeneration have not been identified (RYCHEL & SWALLA, 2009; MIYAMOTO & SAITO, 2010).

LIFE HABITS

Acorn worms are exclusively marine and usually burrow into the sand or mud or are associated with the underside of rocks, although species have also been found entangled in kelp holdfasts (RITTER, 1900). To burrow, they penetrate the sediment using muscular peristaltic contractions of the proboscis and transport the sediment using the ciliated epidermis of the proboscis and collar. Burrow structure varies greatly. They may be Y-shaped, with one arm forming a feeding funnel (DUNCAN, 1987), U-shaped (RAO, 1954), coiled (BURDON-JONES, 1950) (Fig. 4.1), sinuous, irregular (BRAMBEL & GOODHART, 1941; KNIGHT-JONES, 1953), poorly defined (BURDON-JONES, 1956) or absent for animals that occupy interstitial spaces (CAMERON, 2002b; WORSAAE & others, 2012). Most acorn worms form mucus-lined burrows (Fig. 4.1) that smell of haloindols or brominated phenols (HIGA, FUJIYAMA, & SCHEUER, 1980; GIRAY & KING, 1997; WOODIN, WALLA, & LINCOLN, 1987). In the absence of a burrow or in response to stress, a mucous tube may be produced (BURDON-JONES, 1952; HALANYCH & others, 2013). The antibiotic properties of the phenols stabilize the sediment around the burrow (CEDHAGEN & HANSSON, 2013) and likely repel predators, such as gastropods on the coasts of Japan and South Carolina that have been known to nip the tail from defecating ptychoderids (K. TAGAWA, E. E. RUPPERT, personal communications). *Saxipendium* WOODWICK & SENSENBAUGH, 1985 cling to rocky outcroppings near hydrothermal vents (WOODWICK & SENSENBAUGH, 1985), and members of the deep-sea family Torquaratoridae reside on the seafloor.

Enteropneusta are mucociliary feeders, that collect their food particles on the proboscis with abundant mucin and glycoprotein secretions and transport them to




FIG. 4. Enteropneust traces. 1, Helical-shaped burrow of *Saccoglossus bromophenolosus* KING, GIRAY, & KORNFELD, 1994, Willapa Bay, Washington, USA, exposing the brominated lining of the burrow (new, photo, M. K. Gingras); 2, fecal casting of *Saccoglossus pusillus* RITTER, 1902 (new); 3, fecal trail of *Tergivelum* sp? (photo courtesy of NOAA Office of Ocean Exploration and Research); 4, resting trace of torquaratorid enteropneust (Twitchett, 1996, fig. 2.1).

the mouth with cilia (BARRINGTON, 1940; KNIGHT-JONES, 1953; BURDON-JONES, 1962; THOMAS, 1972). Many are facultative filter feeders that pump water into the mouth, then through the pharynx and gills slits using cilia that line the gill bars (BARRINGTON, 1940; KNIGHT-JONES, 1953; BURDON-JONES, 1962; CAMERON, 2002a; GONZALEZ & CAMERON, 2009). Balanoglossids ingest copious amounts of sand, in some cases forming feeding funnels at the mouth of the burrow system. *Saccoglossus* extend a long proboscis from the burrow to exploit surface sediments and marine snow, sometimes forming feeding rosettes. To defecate, the worm protrudes its posterior from the

burrow system and ejects the fecal casting with rapid force. These surface castings are often the best clue that acorn worms reside below the sediment surface (Fig. 4.2).

Torquaratorids have evolved several unique traits for life in the deep sea. Most notably, they are often brightly colored and, rather than residing in burrows, they are epibenthic deposit feeders (Fig. 4.3). The collar of many species is elongated into lips that are used to collect and transport sediment to the mouth (HOLLAND & others, 2005; OSBORN & others, 2012; PRIEDE & others, 2012). Their castings form tightly wound spiral coils to wandering, switchback loops on the sea floor (BOURNE & HEEZEN

1965; SMITH, HOLLAND, & RUHL, 2005) (Fig. 4.3). Trace fossils may provide some indirect evidence of acorn worms in the fossil record. See MALETZ (2014) for a review of trace fossils with possible enteropneust affinities, including ones of a spiral form. Many typical meandering deep-water trace fossils (e.g., graphoglyptid traces, *Nereites* ichnofacies) may be explained as mucus-coated fecal casts produced by enteropneusts. The gut contents act as ballast, and when voided, the worms may drift through the demersal zone (SMITH, HOLLAND, & RUHL, 2005; OSBORN & others, 2012; HOLLAND, KUHNZ, & OSBORN, 2012). This adaptation allows for economical, long-distance movements to a new feeding site in an environment where food is typically heterogeneous and limited (OSBORN & others, 2012; JONES & others, 2013).

Two species of *Glandiceps* SPENGLER, 1893 from the west Pacific swim by flattening the trunk in a dorsoventral aspect, and then undulate and turn as the worm ascends to the sea surface (SPENGLER, 1909; YOSHIMATU & NISHIKAWA, 1999; URATA, IWASAKI, & OHTSUKA, 2012). This behaviour does not appear to be coordinated with spawning (URATA, IWASAKI & OHTSUKA, 2012). An unusual feature of some ptychoderids is their ability to produce light upon stimulation (KUWANO, 1902; CROZIER, 1917; HARVEY, 1926; RAO, 1954). This luminescence is neurally mediated and appears to be intracellular, though luminous slime is also produced (BAXTER & PICKENS, 1964). The role of swimming and bioluminescence in acorn worms is unknown.

EVOLUTION

The embryologist BATESON (1884, 1885) studied the direct-developing acorn worm *Saccoglossus kowalevskii* AGASSIZ, 1873 and found the notochord, gill slits, collar nerve cord, and mesoblast development comparable to amphioxus, and so regarded enteropneusts as sister to chordates. Earlier, METSCHNIKOFF (1869) studied the development of the genus *Balanoglossus* DELLE

CHIAIE, 1829 and noted many similarities of the tornaria to echinoderm larvae, so regarded hemichordates as sister taxon to echinoderms, a superphyletic group that he called Ambulacraria. With the advent of modern cladistics and phylogenetic methods, early phylogenies were consistent with these early ideas that envisioned the deuterostome ancestor as either a sessile tentaculate filter-feeding organism similar to a pterobranch (ROMER, 1967), or to a diplurula larva (GARSTANG, 1928). This dichotomy, and the polyphyletic relationship of enteropneusts (as sister to chordates) and pterobranchs (allied with echinoderms), persisted for nearly a century (GEE, 1996; HOLLAND, HOLLAND, & HOLLAND, 2015; LOWE & others, 2015; SATOH, 2016; PETERSON & EERNISSE, 2016). These hypotheses have fallen out of favor, replaced by the hypothesis that the deuterostome ancestor was a free-living worm with a pharynx perforated with gill slits, used in filter feeding—similar to a modern enteropneust (e.g., CAMERON, SWALLA, & GAREY, 2000; CAMERON, 2002b; LOWE & others, 2015; SATOH, 2016). This hypothesis is consistent with an extensive morphological phylogeny for hemichordates (CAMERON, 2005) and current molecular phylogenies, which regard Enteropneusta and Pterobranchia as sister taxa and the monophyletic Hemichordata as sister to the Echinodermata, together forming the Ambulacraria (Fig. 5) (WINCHELL & others, 2002; PETERSON & others, 2013; CANNON & others, 2014; TELFORD & others, 2014; SIMAKOV & others, 2015).

The class Enteropneusta consists of three major clades, one comprised of the three families Ptychoderidae, Torquaratoridae, and Spengelidae; the monophyletic Harrimaniidae (OSBORN & others, 2012; CANNON & others, 2014); and the stem group Cambrian fossils (Fig. 5). According to a molecular clock estimate based on the genomes of *Ptychodera flava* and *Saccoglossus kowalevskii*, the three-family clade and Harrimaniidae clade diverged 373 million years ago (SIMAKOV & others, 2015). Among




FIG. 5. Phylogenetic tree of acorn worms. The age of each fossil is provided in millions of years, and the stratigraphic details are provided in the text. Neither the tree branch lengths nor the worm silhouettes are to scale. The age of the Ambulacraria, Hemichordata, and crown group are molecular clock estimates from SIMAKOV & others (2015) (adapted from Cameron, 2016).

the larger three-family clade, Spengelidae is sister to the Ptychoderidae + Torquaratoridae (CAMERON, 2005; OSBORN & others, 2012; CANNON & others, 2014). OSBORN and others (2012), sequenced 25 torquaratorids and found them to be a monophyletic sister group to the Ptychoderidae. Based on two unidentified specimens, CANNON and others (2014) suggested that the Torquaratoridae are nested within the family Ptychoderidae (i.e., ptychoderid paraphyly). Torquaratorids have several unique morphological adaptations to life in the deep

sea (see family diagnoses, below) that are absent from ptychoderids. Interestingly, these adaptations did not evolve by parallelism in the deep-sea acorn worms *Glandiceps abyssicola* SPENGLER, 1893 (Spengelidae) (HOLLAND & others, 2013), *Ritteria* DELAND & others, 2010, nor *Saxipendium* (Harrimaniidae) (DELAND & others, 2010; HOLLAND, OSBORN, & KUHNZ 2012).

The family Harrimaniidae is monophyletic and sister to the three-family clade (Fig. 5). Within the Harrimaniidae there are three clades that differ in the arrangements

of their proboscis musculature (DELAND & others, 2010). *Saxipendium*, a genus previously assigned to the monospecific family Saxipendidae, has been reclassified as a harrimaniid based in part on the shared presence of diffusely arranged proboscis musculature with *Mesoglossus* DELAND & others, 2010 and *Ritteria*. The proboscis muscles of *Saccoglossus* are arranged in concentric rings, whereas radially arranged muscles are the ancestral acorn worm state (DELAND & others, 2010).

The three-family clade and Harrimaniidae clade, inclusive of their fossil species, together comprise the crown Enteropneusta and diverged from a common ancestor about 373 million years ago (Fig. 5). Neither clade can be regarded as ancestral or derived. Each of the clades (as well as each family and species) exhibits ancestral and derived traits, though in general, the evolution of harrimaniids is characterized by losses of tornaria larval traits, whereas that of ptychoderids is characterized by gains of adult complexity. Harrimaniidae develop via a swimming, non-feeding, ciliated larva that swims with a telochrochal ciliated band—an acorn worm innovation. The harrimaniid larval telotroch suggests that the tornaria larva was lost in this family.

The modern families are a crown group clade with respect to the Cambrian fossil stem group, which includes *Oesia disjuncta* and *Spartobranchus tenuis* (Fig. 5). These 505 million-year-old fossils then, are the closest representatives of the ancestral acorn worm condition, and in most respects, they have a harrimaniid-like body plan. The Cambrian fossils lack hepatic sacs, genital wings, and synaptaculæ that bridge the primary and secondary gill bars that characterize ptychoderids and the spengelids *Spengelia* WILLEY, 1898 and *Schizocardium*—this trait then, evolved by parallelism in Cephalochordata (Figs. 3.1–3.2). Tubes are unique to stem enteropneusts. Those of *Oesia disjuncta* are fibrous (Fig. 6.1) and those of *Spartobranchus tenuis* sometimes branch (Fig. 6.2). Dozens of the *O. disjuncta* and approximately one-quarter of the *S. tenuis* specimens

are associated with tubes, suggesting a facultative tubicolous habit (CARON, CONWAY MORRIS, & CAMERON, 2013; NANGLU & others, 2016). The tubes are interpreted as a homolog to pterobranch tubes, abandoned on the internode branch to the crown enteropneust clade.

PALEONTOLOGY

Fossils of the soft-bodied enteropneusts are rare and typically occur in Konservat-Lagerstätten. The oldest of these are the Cambrian fossils *Oesia disjuncta* and *Spartobranchus tenuis* from the Burgess Shales of British Columbia, Canada, and date to about 505 million years ago. *O. disjuncta* is abundant in the Raymond Quarry and the Marble Canyon Quarry (Fig. 7.2). Its body is dominated by a broad and long pharynx that extends to nearly the posterior end of the animal. There is no appreciable intestine. A few of the specimens show what appears to be a posterior grasping appendage. Based on these unusual traits, NANGLU and others (2016), speculated that it may be a basal hemichordate, but here it is allied with *S. tenuis* in the stem Enteropneusta, with which it shares tubes and a similar age. *O. disjuncta* occupied tubes that were formerly classified as the algae *Margaretia* WALCOTT, 1931. The tubes are cosmopolitan, composed of woven fibers that form a sediment anchor basally and project into the water column (Fig. 6.1). The large pharynx, tube-dwelling habit, and lack of sediment in the gut suggest it was an epibenthic filter feeder. *S. tenuis*, from the Walcott Quarry, has a body plan that resembles that of a modern harrimaniid worm, including an extensive pharynx followed by a vermiform, unembellished intestine (Fig. 7.1a–c). It occupied infaunal tubes that could tear, though fibers are not apparent. The discovery of a torquaratorid inside a mucous tube has led to the idea that *S. tenuis* was a member of the family Torquaratoridae (HALANYCH & others, 2013; CANNON & others, 2014). This idea is rejected because *S. tenuis* lacks all diagnostic torquaratorid traits, mucus does not fossilize

(NANGLU, CARON, & CAMERON, 2015), and the Cambrian fossils predate the origin of the crown enteropneusts (SIMAKOV & others, 2015). *S. tenuis* was probably a facultative filter and deposit feeder. The tubes of Cambrian acorn worms provide evidence of a link to the tubarium of pterobranchs, including graptolites (CARON, CONWAY MORRIS, & CAMERON, 2013; NANGLU & others, 2016).

Following the Cambrian, over 200 million years passed before the next fossil acorn worms appeared in the Carboniferous, and after the divergence of the modern families, at about 373 million years ago (SIMAKOV & others, 2015). Though a precarious exercise, the Carboniferous, Triassic, and Jurassic acorn worms have been assigned to the four modern families: Harrimaniidae, Spengelidae, Ptychoderidae, and the deep-sea Torquaratoridae (CAMERON, 2016). *Mazoglossus ramsdelli* (see BARDACK, 1985, 1997; MALETZ, 2014) and *Saccoglossus testa* CAMERON, 2016 are from the Mazon Creek fauna, Illinois, USA (~307–309 million years ago). *M. ramsdelli* (see p. 13) has a wide anterior trunk suggesting a well-developed pharynx (Fig. 8,3). Gill pores are reported (BARDACK, 1985), but having viewed all of the specimens to date, the present author was unable to detect them. The enlarged gonadal-branchial region and absence of gonadal wings suggest it was a member of the family Spengelidae (CAMERON, 2016). The two *S. testa* specimens, also from Mazon Creek, are located next to each other in a single concretion (Fig. 8,1). They possess a long proboscis and therefore likely represent a re-emergence to surface deposit feeding. The proboscis of living saccoglossids are aposomatically colored—usually orange, red, or peach—and chemically defended. The Mazon Creek fauna is replete with predators including horseshoe crabs, sea scorpions, crustaceans, a shark, and the Tully monster (MCCOY & others, 2016; SALLAN & others, 2017), so chemical defense and aposomatic colors may have evolved as early as *S. testa* (CAMERON, 2016).




FIG. 6. Tubes of Cambrian fossil acorn worms. 1, spirally arranged pores perforate the tube of *Oesia disjuncta* WALCOTT, 1911, ROM 63716 (Nanglu & others, 2016, fig. 4a); 2, branching tubes of *Spartobranchus tenuis* (WALCOTT, 1911), ROM 57170 (Caron, Conway Morris, & Cameron, 2013, fig. 2f).

The next evidence of an acorn worm is a resting trace fossil from the Lower Triassic Werfen Formation of northern Italy (TWITCHETT, 1996) (Fig. 4.4). It has collar lips that extend laterally and a pharynx that narrows posteriorly, characteristic of the Torquaratoridae, a family of surface-dwelling, pelagic deep-sea acorn worms (HOLLAND & others, 2005; PRIEDE & others, 2012). *Megaderiaion sinemuriense* (see p. 13) is from the Sinemurian (Lower Jurassic) strata of Osteno, northern Italy (Fig. 8,2). It is a single, small (2 cm) specimen that lacks gonadal wings, hepatic sacs, and an enlarged branchial region, characteristic of the family Harrimaniidae. It resembles *Mesoglossus*, a harrimaniid worm with a medium-long proboscis (DELAND & others, 2010).

Ptychodera callovianum (ALESSANDRELLO, BRACCHI, & RIOU, 2004) was a ptychoderid (Fig. 8,4), from the Lower Callovian (Middle Jurassic) of La Voulte-sur-Rhone, France. The youngest fossil is *Mesobalanoglossus buergeri* (see p. 14) from the lower Tithonian (Upper Jurassic) Solnhofen Limestones, southern Germany (Fig. 8,5). It is assigned to Ptychoderidae due to its length (68.8 cm) and enlarged genital wings, but the proboscis

preservation is poor and the presence of bristles in the pharynx (FRICKINGER, 1999) suggest that it was not an acorn worm.

This chapter limits its systematic section to seven genera with fossil evidence—two from the stem group, and five from the crown group. The excluded genera are only known from extant species. What we understand at this point, based on the pattern of appearance of these seven fossil acorn worm species, is: (1) the evolution of acorn worms is characterized by an evolutionary stasis—the harrimaniid body plan of *Spartobranchus tenuis* appears in the Cambrian and persists to this day in the Harrimaniidae, including *Saccoglossus* (Fig. 5); (2) this body plan was followed by that of a spengelid in *Mazoglossus ramsdelli* from the Pennsylvanian (BARDACK, 1997) and then by a torquaratorid, characterized by collar lips, in a Lower Triassic trace fossil (TWITCHETT, 1996); and (3) the complex ptychoderid body plan appeared last in *Ptychodera callovianum* from the Upper Jurassic (ALESSANDRELLO, BRACCHI, & RIOU, 2004; CAMERON, 2016) (Fig. 5).

STEM GROUP TAXA

The stem group enteropneusts include tube-dwelling taxa with a vermiform body that consists of a proboscis, a collar, and a trunk with a pharynx framed by multiple, paired, circumferential gill bars and slits.

Oesia WALCOTT, 1911, p. 132 [**O. disjuncta*; OD]

Tripartite body with proboscis, collar, and trunk; trunk unusual, possessing extensive pharynx and lacking esophagus and intestine; posterior grasping appendage; tubes, fibrous with spirally arranged openings. *Cambrian (Miaolingian, Wuliuan, Bathyriscus–Elrathia Biozones)*: Canada.—FIG. 7.2. **O. disjuncta*, USNM 57630, lectotype (selected by SZANIAWSKI, 2005, p. 2), one of three specimens illustrated by WALCOTT (1911), Burgess Shale, Walcott Quarry, British Columbia, Canada (new, photo by J.-B. Caron).

Spartobranchus CARON, CONWAY MORRIS, & CAMERON, 2013, p. 503 [**Ottoia tenuis* WALCOTT, 1911, p. 130; OD]. Vermiform, maximum length 10 cm; body comprised of short proboscis, collar, pharyngeal area with up to 40 pairs of gill and tongue bars; elongate posterior trunk ending in bulbous unit; gut straight, anus terminal; approximately one-quarter of specimens associated with tube of organic walls; tube varies from straight, completely

circular, to helicoidal, and sometimes branching; no more than one individual per tube has been observed. *Cambrian (Miaolingian, Wuliuan, Bathyriscus–Elrathia Biozones)*: Canada.—FIG. 7.1*a–c*. **S. tenuis* (WALCOTT); 1*a–b*, USNM 108494, lectotype (selected by Caron, Conway Morris, & Cameron, 2013, supplement 1, from a small number of specimens supposedly collected by WALCOTT [1911] who described, but never illustrated *Ottoia tenuis* and the whereabouts of his material is unknown), specimen missing proboscis and terminal portion of trunk; 1*a*, part, direct light; 1*b*, counterpart, polarized light; 1*c*, complete specimen, ROM 62123 (Caron, Conway Morris, & Cameron, 2013, fig. 1).

CROWN GROUP TAXA

Family HARRIMANIIDAE

Spengel, 1901

[Harrimaniidae SPENGLER, 1901, p. 215]

Simple morphology, largely characterized by features it lacks; no dorsal collar nerve roots arising from the collar nerve cord; gills lack synaptaeulae that bridge primary and secondary gill bars; trunk lacks hepatic sacs, circular muscle fibers, and lateral septa, only dorsal and ventral septa present; intestinal pores only rarely present; posterior projecting horns of Y-shaped proboscis skeleton extend at least to middle of collar; proboscis longitudinal musculature may be arranged diffusely, in radial plates, or in concentric rings; development via short-lived, non-feeding larvae that become juveniles with adhesive post-anal tail. *Carboniferous (Pennsylvanian, Moscovian)–Holocene* (extant): worldwide.

The Harrimaniidae comprises ten living genera, including one (*Saxipendium* WOODWICK & SENSENBAUGH, 1985) that was previously assigned to its own family (DELAND & others, 2010; WORSAAE & others, 2012). *Saccoglossus* is the most speciose and familiar genus (CAMERON, DELAND, & BULLOCK, 2010). Its proboscis longitudinal musculature is arranged in concentric rings, while other genera may be arranged diffusely (e.g., *Mesoglossus*), or in radial plates (e.g., *Protoglossus* HORST, 1939).

Saccoglossus SCHIMKEWITSCH, 1892, p. 2 [**Balanoglossus mereschkowskii* WAGNER, 1885, p. 46; OD] [= *Balanoglossus (Dolichoglossus)* SPENGLER, 1893,




FIG. 7. Stem group Enteropneusta (p. 12).

p. 360 (type, *D. kowalewskii* AGASSIZ, 1873, SD SPENGLER, 1901, p. 215)]. Proboscis long; longitudinal muscle fibers of proboscis arranged in several concentric rings; middorsal longitudinal groove may be present; collar usually very short compared to proboscis; dorsal interbranchial genital ridges and dorsal gonads absent but lateral extrabranchial genital ridges may be present; intestinal pores often present; perihæmal cavities always present; peribuccal cavities usually present. [Many species favor quiet muddy-sandy flats not far from the mouth of a bay, living in semi-permanent and helical-shaped burrows and throwing up low conical mounds of quasi-spiral fecal castings. See CAMERON, DELAND, & BULLOCK, 2010; also see Fig 4.2.]—FIG. 8, 1. *S. testa* CAMERON, 2016, holotype, FMNH PE 45216, one of two counterparts (new).

Megaderaion ARDUINI, PINNA, & TERUZZI, 1981, p. 105 [*M. sinemuriense*; OD]. Elongate worm divided into elongate, rounded, and ogival proboscis; collar wider than long, tapering trunk; single specimen small (2 cm) lacking gonadal wings, hepatic sacs, and enlarged branchial region, characteristic of family Harrimaniidae; resembles *Mesoglossus*, with medium-long proboscis. [See DELAND & others, 2010.] *Lower Jurassic (Sinemurian, Coroniceris bucklandi Biozone)*: Italy.—FIG. 8, 2. *M. sinemuriense*, holotype, i751 (Museo Civico di Storia Naturale di Milano) (Arduini, Pinna, & Teruzzi, 1981, fig. 1).

Family SPENGLIDAE Willey, 1899

[Spengelidae WILLEY, 1899, p. 239] [=Glandicipitidae SPENGLER, 1901, p. 215]

Anterior vermiform process of stomochord present in all members; skeletal horns usually extend over whole length of collar; dorsal

nerve roots arising from collar nerve cord rare; lateral septum absent; hepatic caeca and synapticula may or may not be present; circular muscle fiber layer positioned inside longitudinal muscle layer in trunk. *Carboniferous (Pennsylvanian, Moscovian)–Holocene* (extant); worldwide.

In the cases where development has been documented, Spengelidae species have a tornaria larva (CAMERON & PEREZ, 2012). The family includes four living genera (see HORST, 1939; CAMERON & PEREZ, 2012). CAMERON (2016) referred *Mazoglossus ramsdelli* to the family Spengelidae, thus, extending an extant enteropneust family to include a Carboniferous fossil species.

Mazoglossus BARDACK, 1997, p. 89 [*M. ramsdelli*; OD] Small (specimens less than 10 cm total length); recognizable as an enteropneust by outline shape of proboscis, collar, and trunk. Illinois, USA.—FIG. 8, 3. *M. ramsdelli*, FMNH PE 23053, one of two counterparts, complete specimen (adapted from Cameron, 2016, fig. 2C).

Family PTYCHODERIDAE Spengel, 1893

[Ptychoderidae SPENGLER, 1893, p. 359] [=Balanoglossidae WILLEY, 1899, p. 239]

Lateral septa in trunk; invariable absence of abdominal pores; lack of vermiform process of stomochord; dorsal nerve roots

in collar present; skeletal horns rarely reach beyond anterior half of collar; synapticulae form bridges between primary and secondary gill bars in pharynx, hepatic caeca usually present; dorsolateral ciliated grooves in abdominal part of alimentary canal; circular muscle fibers in trunk, usually outside the longitudinal fibers; development occurs through typical tornaria larva [See CAMERON & OSTIGUY, 2013, p. 144.] *Middle Jurassic (Callovian)–Holocene* (extant): worldwide.

Ptychoderidae includes four extant genera, including the familiar *Balanoglossus*. *Mesobalanoglossus* is here included in the Ptychoderidae. CAMERON (2016) referred the Upper Jurassic *Megaderaion callovanum* ALESSANDERELLO, BRACCHI, & RIOU, 2004 to the Ptychoderidae. The species is here included in the genus *Ptychodera*.

Ptychodera ESCHSCHOLTZ, 1825, p. 740 [**P. flava*; M]. As for family, except gill pores open by long slits, whereas those of *Balanoglossus* open by small pores.—FIG. 8,4. *P. callovanum* (ALESSANDERELLO, BRACCHI & RIOU, 2004), MNHN L.P.M.-B. 48352, holotype.

Mesobalanoglossus BECHLY & FRICKHINGER in FRICKHINGER, 1999, p. 77 [**M. buergeri*; OD] Single specimen, 68.8 centimeters long, with diminutive proboscis, collar, and long trunk with bristles interpreted as gill bars; trunk tapers posteriorly, anterior wide part interpreted as gonadal wings. *Upper Jurassic (Lower Tithonian, Hybonoticeris hybonotum Biozone)*: Germany. —FIG. 8,5. **M. buergeri*, holotype (Bechly & Frickhinger in Frickhinger, 1999, fig. 144).

Family TORQUARATORIDAE Holland & others, 2005

[Torquaratoridae HOLLAND & others, 2005, p. 374]

Proboscis broad, short, dome shaped; collar wide, usually embellished with lateral lips; body semi-transparent, frequently brightly colored; muscular system poorly developed; collagenous proboscis skeleton and gill bars greatly reduced; proboscis skeletal horns absent from all species but *Torquarator bullocki* (HOLLAND & others, 2005). *Lower Triassic (upper Olenekian)–Holocene* (extant): worldwide.

The family Torquaratoridae is exclusive to the deep sea (HOLLAND & others, 2005, 2009; HOLLAND, KUHNZ, & OSBORN, 2012).

There are five genera (JABR, ARCHAMBAULT, & CAMERON, in press). Fossil specimens of the Torquaratoridae are unknown, but a single resting trace from the Lower Triassic (Olenekian) Werfen Formation of northern Italy (TWITCHETT, 1996) shows, through its characteristic shape, the presence of the family in the Mesozoic (see Fig. 4.4).

FUTURE RESEARCH

In the intervening years since the first edition of this chapter (BULMAN, 1970) in which only a short note on the Enteropneusta appeared, 22 new, living enteropneust species have been described, and with it, their zoogeographic range better appreciated (see especially, CAMERON, DELAND, & BULLOCK, 2010), particularly with respect to their abundance in the deep sea (OSBORN & others, 2012; PRIED & others, 2012; CANNON, SWALLA, & HALANYCH, 2013). While fossil enteropneusts were unknown to BULMAN (1970), here we show that seven fossil acorn worm species extend from the Jurassic to the Cambrian periods. Fine details of cellular ultrastructure (see especially PARDOS & BENITO, 1990) have been brought into the light, contributing to a better appreciation of the functional morphology of the group. Surprising observations have been made on their life habits. Interest in the molecular developmental of the group has revealed body plan patterns conserved with their common ancestor, chordates (LOWE & others, 2003; LOWE & others, 2006). The current phylogeny is robust, with broad taxonomic sampling (CANNON & others, 2014), with the exception of the family Spengelidae. The genomes of *Saccoglossus kowalevskii* and *Ptychodera flava* have been sequenced, providing a tremendous resource for future investigations (TAGAWA, 2016; SIMAKOV & others, 2015). The conserved linkage, or micro-synteny, of transcription factors involved in gill development have been revealed (GILLIS, FRITZENWANKER, & LOWE 2012; SIMAKOV & others, 2015). Similar studies on other organ systems will come, and higher levels of conserved linkages, or




FIG. 8. Harrimaniidae (1–2), Spengelidae (3), Ptychoderidae (4–5). (p. 12–14).

macro-synteny, may be expected. The genes involved in the biosynthesis and degradation of the sialic acid molecule, a deuterostome synapomorphy (CAMERON, 2005), includes nine that arrived as transposons from a prokaryote or protist donor (SIMAKOV & others, 2015). In their review of the development of the group, KAUL-STREHLOW and RÖTTINGER (2015) stressed a need for more investigations on the development of the nervous and muscular systems, and I would add to that the coelomic, reproductive, and immune systems. There

is a single paper on the immunodefense of the group (TASSIA, WHELAN, & HALANYCH, 2017). Interest in acorn worm biology has grown—affirmed by the 1st International Hemichordate Meeting, Hopkins Marine Station at Stanford University, December 2016—and the future looks bright.

Future fruitful avenues of research may include: (1) A more extensive sampling of South America, Africa, Asia, as well as deep water where many species remain to be found. (2) The burrow shape of acorn worms varies tremendously and there seems

to be some genus-specific patterns. The fossil record of burrow trace fossils is rich (BROMLEY, 1996; SEILACHER, 2007; GINGRAS & others, 2008), and may provide some insight on the origin of modern acorn worm genera. HORST (1940) suggested that an enteropneust similar to *Saccoglossus* is responsible for the spiraling trace fossil, *Gyrolithes*. This, as well as changes to burrow structure—depending on season or sediment parameters—should be explored. (3) Intensive sampling of many individuals within a species is needed to assess the amount of interspecific variation, in the structure of the heart-kidney coelomic process, the collagenous gill bars and nuchal skeleton, the ossicles, as well as developmental genes—since variation is what selection works on. Such studies are rare, the most comprehensive study being that of the many *Psychodera flava* variants from the Indian Ocean (PUNNETT, 1906). (4) Do closely related species have similar developmental processes—as might be expected—or is the variation substantive, as we have seen from the direct developer *Saccoglossus kowalevskii*, to the tornaria of *Schizocardium californicum* (see GONZALEZ, UHLINGER, & LOWE, 2017)? Experimental molecular developmental approaches that include gene knock-down (TAGAWA, 2016), RNAi, CRISPR, and single-cell sequencing (KLEIN & others, 2015), will provide more robust and nuanced tests of homology. (5) Nearly nothing is known of the post-juvenile development of acorn worms, but the powerful potential of these investigations is that the ontological development can be compared with the paleontological equivalents, particularly with the Cambrian forms that are abundant and show exquisite details of extracellular matrix structures, including gill bars and tubes. (6) Nothing is written on the quantitative genetics, the population-based studies of continuously varying characters, nor how these characters respond to selection. (7) Slight more is known about the neural physiology (PICKENS, 1970; CAMERON & MACKIE, 1996), and respiratory physiology of acorn worms (AZARIAH, ISMAIL, &

NAJIB, 1975; DITADI, MENDES, & BIANCONCINI, 1997). Is the physiological role of the gill slits respiratory as the name Enteropneusta—Greek for gut-breathing—implies, or is the primary role of the acorn worm gills in ammonia exchange or acid-base regulation? Even less is known of embryonic and larval physiology and behavior.

ABBREVIATIONS FOR MUSEUM REPOSITORIES

BSPGM: Bayerische Staatssammlung für Paläontologie und Historische Geologie, Munich, Germany
FMNH PE: Field Museum, Chicago, Illinois, USA
i751: Museo Civico di Storia Naturale di Milano, Italy
ROM: Royal Ontario Museum, Toronto, Canada
USNM: United States National Museum of Natural History, Washington, D. C., USA
MNHN: Muséum National d'Histoire Naturelle, Paris, France

REFERENCES

- Agassiz, Alexander. 1873. The history of *Balanoglossus* and Tornaria. *Memoirs of the American Academy of Arts & Sciences* (new series) 9:421–436.
- Alessandrello, Anna, Giacomo Bracchi, & Bernard Riou. 2004. Polychaete, sipunculan and enteropneust worms from the Lower Callovian (Middle Jurassic) of La Voulte-sur-Rhône (Ardeche, France). *Memorie della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano* 32(1):1–16.
- Arduini, Paolo, Giovanni Pinna, & Giorgio Teruzzi. 1981. *Megaderiaion sinemuriense*, n. g., n. sp., a new fossil enteropneust of the Sinemurian of Osteno in Lombardy. *Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano* 122:104–108.
- Azariah, Jayapaul, M. Mohamed Ismail, & M. Ahmed Najib. 1975. Investigation on the ecology and respiratory responses of the hemichordate *Psychodera flava* to tidal cycles and salinity changes. *The Biological Bulletin* 149(3):455–466.
- Balsler, E. J., & E. E. Ruppert. 1990. Structure, ultrastructure, and function of the preoral heart-kidney in *Saccoglossus kowalevskii* (Hemichordata, Enteropneusta) including new data on the stomochord. *Acta Zoologica (Stockholm)* 71:235–249.
- Bardack, David. 1985. Les premiers fossiles de hagfish (Myxiniiformes) et Enteropneusta (Hemichordata) dépôts de la faune (Pennsylvanienne) du Mazon Creek dans l'Illinois, U.S.A. *Bulletin trimestriel de la Société d'histoire naturelle et des amis du Muséum d'Autun* 116:97–99.
- Bardack, David. 1997. Wormlike Animals: Enteropneusta (acorn worms). In C. W. Shabica & A. A. Hay, eds., *Richardson's Guide to the Fossils of the Mazon Creek*. Northeastern Illinois University, Chicago. p. 89–92.

- Barrington, E. J. W. 1940. Observations on feeding and digestion in *Glossobalanus minutus*. Quarterly Journal of Microscopical Science 82:227–260.
- Bateson, William. 1884. The early stages in the development of *Balanoglossus*. Quarterly Journal of Microscopical Science 24:208–236.
- Bateson, William. 1885. The later stages in the development of *Balanoglossus kowalevskii*, with a suggestion as to the affinities of the Enteropneusta. Quarterly Journal of Microscopical Science (Supplement) 25:81–122.
- Baxter, C. H., & P. E. Pickens. 1964. Control of luminescence in hemichordates and some properties of a nerve net system. Journal of Experimental Biology 41:1–14.
- Benito, Jesús, & Fernando Pardos. 1997. Hemichordata. In F. W. Harrison & E. E. Ruppert, eds., Microscopic Anatomy of Invertebrates. Vol. 15. Wiley-Liss. New York. p. 15–101.
- Bourne, D. W., & B. C. Heezen. 1965. A wandering enteropneust from the abyssal Pacific, and the distribution of “spiral” tracks on the sea floor. Science (new series) 150(3692):60–63.
- Brambell, F. W. R., & H. A. Cole. 1939. The preoral ciliary organ of the Enteropneusta: Its occurrence, structure, and possible phylogenetic significance. Proceedings of the Zoological Society, London 109:181–193.
- Brambell, F. W. R., & C. B. Goodhart. 1941. *Saccoglossus horsti* sp. n., an enteropneust occurring in the Solent. Journal of the Marine Biological Association of the United Kingdom 24:283–301.
- Bromley, R. G. 1996. Trace Fossils. Biology, Taphonomy and Applications. Chapman & Hall. London. 361 p.
- Bromley, R. G. 2012. Trace Fossils: Biology, Taxonomy and Applications. Unwin Hyman. London. 280 p.
- Bullock, T. H. 1940. The functional organization of the nervous system of Enteropneusta. Biological Bulletin 79:91–113.
- Bulman, O. M. B. 1970. Graptolithina. In Curt Teichert, ed., Treatise on Invertebrate Paleontology, Part V, Second Edition. The Geological Society of America & The University of Kansas Press. Boulder & Lawrence. xxxii + 163 p.
- Burdon-Jones, Cyril. 1950. An enteropneust genus new to the British Isles. Nature 3165:327.
- Burdon-Jones, Cyril. 1951. Observations on the spawning behaviour of *Saccoglossus horsti* Brambell & Goodhart, and of other Enteropneusta. Journal of the Marine Biological Association of the United Kingdom 29:625–638.
- Burdon-Jones, Cyril. 1952. Development and biology of the larva of *Saccoglossus horsti* (Enteropneusta). Philosophical Transactions of the Royal Society of London (series B) 236:553–589.
- Burdon-Jones, Cyril. 1956. Observations on the enteropneust, *Protoglossus koehleri* (Caullery & Mesnil). Journal of Zoology 127:35–58.
- Burdon-Jones, Cyril. 1962. The feeding mechanism of *Balanoglossus gigas*. Boletim da Faculdade de Filosofia, Ciências e Letras, Universidade de São Paulo. Zoologia 24:255–280.
- Cameron, C. B. 2002a. Particle retention and flow in the pharynx of the enteropneust worm *Harrimania planktophilus*: The filter feeding pharynx may have evolved prior to the chordates. Biological Bulletin 202:192–200.
- Cameron, C. B. 2002b. The anatomy, life habits and later development of a new species of enteropneust, *Harrimania planktophilus* (Hemichordata: Harrimaniidae) from Barkley Sound. Biological Bulletin 202:182–191.
- Cameron, C. B. 2005. A phylogeny of the hemichordates based on morphological characters. Canadian Journal of Zoology 83(1):196–215.
- Cameron, C. B. 2016. *Saccoglossus testa* from the Mazon Creek fauna (Pennsylvanian of Illinois) and the evolution of acorn worms (Enteropneusta: Hemichordata). Palaeontology 59(3):329–336, doi:10.1111/pala.12235.
- Cameron, C. B., & C. D. Bishop. 2012. Biomineral ultrastructure, elemental constitution and genomic analysis of biomineralization-related proteins in hemichordates. Proceedings of the Royal Society of London (series B) 279(1740):3041–3048.
- Cameron, C. B., Carine Deland, & T. H. Bullock. 2010. A revision of the genus *Saccoglossus* (Hemichordata: Enteropneusta: Harrimaniidae) with taxonomic descriptions of five new species from the Eastern Pacific. Zootaxa 2483:1–22.
- Cameron, C. B., & G. O. Mackie. 1996. Conduction-pathways in the nervous system of *Saccoglossus* sp. (Enteropneusta). Canadian Journal of Zoology 74:15–19.
- Cameron, C. B., & Angélica Ostiguy. 2013. Three new species of *Glossobalanus* (Hemichordata: Enteropneusta: Ptychoderidae) from western North America. Zootaxa 3630(1):143–154.
- Cameron, C. B., & Maëva Perez. 2012. Spengelidae (Hemichordata: Enteropneusta) from the eastern Pacific including a new species, *Schizocardium californicum*, from California. Zootaxa 3569: 79–88.
- Cameron, C. B., B. J. Swalla, & J. R. Garey. 2000. Evolution of the chordate body plan: New insights from phylogenetic analysis of deuterostome phyla. Proceedings of the National Academy of Sciences 97(9):4469–4474.
- Cannon, J. T., K. M. Kocot, D. S. Waits, D. A. Weese, B. J. Swalla, S. R. Santos, & K. M. Halanych. 2014. Phylogenomic resolution of the hemichordate and echinoderm clade. Current Biology 24:2827–2832.
- Cannon, J. T., B. J. Swalla, & K. M. Halanych. 2013. Hemichordate molecular phylogeny reveals a novel cold-water clade of harrimaniid acorn worms. Biological Bulletin 225:194–204.
- Caron, J.-B., Simon Conway Morris, & C. B. Cameron. 2013. Tubicolous enteropneusts from the Cambrian. Nature 7442:503–506.
- Caullery, Maurice, & Félix Mesnil. 1900. Sur une nouvelle espèce de *Balanoglossus* (*B. koehleri*) habitant les zones de la Manche. Comptes rendus hebdomadaires des Séances et Mémoires de la Société de Biologie 52:256–259.

- Cedhagen, Tomas, & H. G. Hansson. 2013. Biology and distribution of hemichordates (Enteropneusta) with emphasis on Harrimaniidae and description of *Protoglossus bocki* sp. nov. from Scandinavia. *Helgoland Marine Research* 67:251–265.
- Colwin, A. L., & L. H. Colwin. 1953. The normal embryology of *Saccoglossus kowalevskii* (Enteropneusta). *Journal of Morphology* 92:401–453.
- Colwin, A. L., & L. H. Colwin. 1954. Fertilization changes in the membranes and cortical granular layer of the egg of *Saccoglossus kowalevskii* (Enteropneusta). *Journal of Morphology* 95:1–46.
- Crawford, B. J., & F. S. Chia. 1978. Coelomic pouch formation in the starfish *Pisaster ochraceus* (Echinodermata: Asteroidea). *Journal of Morphology* 157:99–120.
- Crozier, W. J. 1917. Photonic sensitivity of *Balanoglossus*. *Journal of Experimental Zoology* 24(2):211–217.
- Dawydoff, Constantin. 1909. Beobachtungen über den Regenerationsprozeß bei den Enteropneusten. *Zeitschrift für wissenschaftliche Zoologie* 93: 237–305.
- Dawydoff, Constantin. 1948. Classe des Entéropneustes. In P. P. Grassé, ed., *Traité de Zoologie*. Masson. Paris. p. 369–453.
- Deland, Carine, C. B. Cameron, T. H. Bullock, K. P. Rao, & W. E. Ritter. 2010. A taxonomic revision of the family Harrimaniidae (Hemichordata: Enteropneusta) with descriptions of seven species from the eastern Pacific. *Zootaxa* 2408:1–30.
- Delle Chiaje, Stefano. 1829. Memorie sulla storia e notomia degli animali senza vertebre del Regno di Napoli 4:1–214.
- Ditadi, A. S. F., E. G. Mendes, & M. S. C. Bianconcini. 1997. Influence of body mass and environmental oxygen tension on the oxygen consumption rates of an enteropneust, *Glossobalanus crozieri*. *Brazilian Journal of Medical and Biological Research* 30(12):1441–1444.
- Duncan, P. B. 1987. Burrow structure and burrowing activity of the funnel-feeding enteropneust *Balanoglossus aurantiacus* in Bogue Sound, North Carolina, USA. *Marine Ecology* 8(1):75–95.
- Eschscholtz, Friedrich. 1825. Bericht über die zoologische Ausbeute der Reise von Kronstadt bis St.-Peter und Paul. Oken's Isis 733–747.
- Ezhova, O. V., & V. V. Malakhov. 2009. Three-dimensional structure of the skeleton and buccal diverticulum of an acorn worm *Saccoglossus mereschkowskii* Wagner, 1885 (Hemichordata: Enteropneusta). *Invertebrate Zoology* 6(2):103–116.
- Frickinger, K. A. 1999. Die Fossilien von Solnhofen 2 [The Fossils of Solnhofen 2]. Goldschneck Verlag. Weinstadt, Baden-Württemberg. p. 76–79.
- Garstang, Walter. 1928. The morphology of the Tunicata and its bearing on the phylogeny of the Chordata. *Quarterly Journal Microscopical Science* 72:51–187.
- Gee, Henry. 1996. Before the Backbone: Views on the Origin of the Vertebrates. Chapman & Hall. London. 346 p.
- Gegenbaur, Carl. 1870. Grundzüge der vergleichenden Anatomie. Zweite, umgearbeitete Auflage. Wilhelm Engelmann. Leipzig. 892 p.
- Gemmill, J. F. 1914. The development and certain points in the adult structure of the starfish *Asterias rubens*. *Philosophical Transactions of the Royal Society of London (series B)* 205:213–294.
- Gilchrist, J. D. F. 1923. A form of dimorphism and asexual reproduction in *Psychodera capensis* (Hemichordata). *Journal of the Linnean Society, London* 35:393–398.
- Gilchrist, J. D. F. 1925. *Xenopleura vivipara*, g. et sp. n. (Enteropneusta). *Quarterly Journal of Microscopical Science* 69:555–573.
- Gillis, J. A., J. H. Fritzenwanker, & C. J. Lowe. 2012. A stem-deuterostome origin of the vertebrate pharyngeal transcriptional network. *Proceedings of the Royal Society of London (series B)* 279:237–246.
- Gingras, M. K., S. E. Dashtgard, J. A. MacEachern, & S. G. Pemberton. 2008. Biology of shallow marine ichnology: A modern perspective. *Aquatic Biology* 2:255–268.
- Giray, Cem, & G. M. King. 1996. *Protoglossus graveolens*, a new hemichordate (Hemichordata: Enteropneusta: Harrimaniidae) from the northwest Atlantic. *Proceedings of the Biological Society of Washington* 109(3):430–445.
- Giray, Cem, & G. M. King. 1997. Predator deterrence and 2,4-dibromophenol conservation by the enteropneusts *Saccoglossus bromophenolosus* and *Protoglossus graveolens*. *Marine Ecology Progress Series* 159:239–238.
- Gislén, T. R. E. 1930. Affinities between the Echinodermata, Enteropneusta and Chordonia. *Zoologiska Bidrag fran Uppsala* 12:199–304.
- Gonzalez, Paul, & C. B. Cameron. 2009. The gill slits and pre-oral ciliary organ of *Protoglossus* (Hemichordata: Enteropneusta) are filter-feeding structures. *Biological Journal of the Linnean Society* 98:898–906.
- Gonzalez, Paul, K. R. Uhlinger, & C. J. Lowe. 2017. The adult body plan of indirect developing hemichordates develops by adding a Hox-patterned trunk to an anterior larval territory. *Current Biology* 27:1–9.
- Hadfield, M. G. 1975. Hemichordata. In A. C. Giese & J. S. Pearse, eds., *Reproduction of Marine Invertebrates*. Academic Press. New York. p. 185–240.
- Haeckel, Ernst. 1879. *Natürliche Schöpfungsgeschichte*. Gemeinverständliche wissenschaftliche Vorträge über die Entwicklungslehre im Allgemeinen und diejenige von Darwin, Goethe und Lamarck im Besonderen. Siebente, umgearbeitete und vermehrte Auflage. Verlag G. Reimer. Berlin. 718 p.
- Halanych, K. M., J. T. Cannon, A. R. Mahon, B. J. Swalla, & C. R. Smith. 2013. Modern Antarctic acorn worms form tubes. *Nature Communications* 4:2738.
- Harvey, E. N. 1926. On the inhibition of animal luminescence by light. *The Biological Bulletin* 51:85–88.
- Henry, J. Q., Kunifumi Tagawa, & M. Q. Martindale. 2001. Deuterostome evolution: Early development in the enteropneust hemichordate, *Psychodera flava*. *Evolution & Development* 3:375–390.
- Higa, Tatsuo, Toraya Fujiyama, & P. J. Scheuer. 1980. Halogenated phenol and indole constituents of acorn worms. *Comparative Biochemistry and Physiology* 65(3):525–530.

- Holland, N. D., D. A. Clague, D. P. Gordon, A. Gebruk, D. L. Pawson, & Michael Vecchione. 2005. 'Lophenteropneust' hypothesis refuted by collection and photos of new deep-sea hemichordates. *Nature* 434:374–376.
- Holland, N. D., L. Z. Holland, & P. W. H. Holland. 2015. Scenarios for the making of vertebrates. *Nature* 520:450–455.
- Holland, N. D., W. J. Jones, Jacob Ellena, H. A. Ruhl, & K. L. Smith, Jr. 2009. A new deep-sea species of epibenthic acorn worm (Hemichordata, Enteropneusta). *Zoosystema* 31(2):333–346.
- Holland, N. D., L. A. Kuhnz, & K. J. Osborn. 2012. Morphology of a new deep-sea acorn worm (class Enteropneusta, phylum Hemichordata): A part-time demersal drifter with externalized ovaries. *Journal of Morphology* 273(7):661–671.
- Holland, N. D., K. J. Osborn, A. V. Gebruk, & Antonina Rogacheva. 2013. Rediscovery and augmented description of the HMS 'Challenger' acorn worm (Hemichordata, Enteropneusta), *Glandiceps abyssicola*, in the equatorial Atlantic abyss. *Journal of the Marine Biological Association of the United Kingdom* 93(8):2197–2205.
- Holland, N. D., K. J. Osborn, & L. A. Kuhnz. 2012. A new deep-sea species of harrimaniid enteropneust (Hemichordata) Proceedings of the Biological Society of Washington 125(3):228–240.
- Horst, C. J. van der. 1934. Die Enteropneusten aus den Meeren der U.S.S.R. *Zoologische Jahrbücher, Abteilung für Anatomie und Ontogenie der Tiere* 58(1):61–81.
- Horst, C. J. van der. 1939. Hemichordata. In H. G. Bronns, ed., *Klassen und Ordnungen des Tier-Reichs wissenschaftlich dargestellt in Wort und Bild*. Akademische Verlagsgesellschaft. Leipzig. Band 4, Abteilung 4, Buch 2, Teil 2. 737 p.
- Horst, C. J. van der. 1940. The Enteropneusta from Inyack Island, Delagou Bay. *South African Museum Annales* 32:293–380.
- Humphreys, Tom, Akane Sasaki, Gene Uenishi, Kekoa Taparra, Asuka Arimoto, & Kuni Tagawa. 2010. Regeneration in the hemichordate *Ptychodera flava*. *Zoological Science* 27:91–95.
- Hyman, L. H. 1959. *The Invertebrates: Smaller Coelomate Groups*, Vol. 5: Chapter 17. The Enterocoelous Coelomates: Phylum Hemichordata, McGraw-Hill. New York, London, & Toronto. p. 72–207.
- Jabr, Noura, Philippe Archambault, & C. B. Cameron. 2018. Biogeography and adaptations of torquaratorid acorn worms (Hemichordata: Enteropneusta) including two new species from the Canadian Arctic. *Canadian Journal of Zoology* [doi 10.1139/cjz-2017-0214].
- Jones, D. O., C. H. Alt, I. G. Priede, W. D. Reid, B. D. Wigham, D. S. Billett, A. V. Gebruk, Antonina Rogacheva, & A. J. Gooday. 2013. Deep-sea surface-dwelling enteropneusts from the Mid-Atlantic Ridge: Their ecology, distribution and mode of life. *Deep Sea Research Part II: Topical Studies in Oceanography* 98:374–387.
- Kaul-Strehlow, Sabrina, & Eric Röttinger. 2015. Hemichordata. In *Evolutionary Developmental Biology of Invertebrates* 6. Springer. Vienna. p. 59–89.
- Kaul-Strehlow, Sabrina, & Thomas Stach. 2013. A detailed description of the development of the hemichordate *Saccoglossus kowalevskii* using SEM, TEM, histology and 3D-reconstructions. *Frontiers in Zoology* 10:53. [doi:10.1186/1742-9994-10-53]
- Kaul-Strehlow, Sabrina, Makoto Urata, Takuya Minokawa, Thomas Stach, & Andreas Wanninger. 2015. Neurogenesis in directly and indirectly developing enteropneusts: of nets and cords. *Organisms Diversity & Evolution* 15:405–422.
- King, G. M., Cem Giray, & Irv Kornfield. 1994. A new hemichordate, *Saccoglossus bromophenolosus* (Hemichordata: Enteropneusta: Harrimaniidae) from North America. *Proceedings of the Biological Society of Washington* 107:383–390.
- Klein, A. M., Linas Mazutis, Ilke Akartuna, Naren Tallapragada, Adrian Veres, Victor Li, Leonid Peshkin, D. A. Weitz, & M. W. Kirschner. 2015. Droplet barcoding for single-cell transcriptomics applied to embryonic stem cells. *Cell* 161:1187–1201.
- Knight-Jones, E. W. 1953. Feeding in *Saccoglossus* (Enteropneusta). *Proceedings of the Zoological Society, London* 123:637–658.
- Kuwano, Hisato. 1902. On a new Enteropneust from Misaki, *Balanoglossus misakiensis* n. sp. *Annotationes zoologicae Japonenses* 4:77–84.
- Lacalli, T. C., & T. H. J. Gilmour. 2001. Locomotory and feeding effectors of the tornaria larva of *Balanoglossus biminienis*. *Acta Zoologica* 82(2): 117–126.
- Lester, S. M. 1985. *Cephalodiscus* sp. (Hemichordata: Pterobranchia): observations of functional morphology, behavior and occurrence in shallow water around Bermuda. *Marine Biology* 85:263–268.
- Linnaeus, Carolus. 1758. *Systema naturae*, per genera tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Tomus I. Editio decima, reformata. Laurentius Salvius. Holmiae. 842 p. [reader.digital-sammlung.de/resolve/display/bsb10076014.html] [www.biodiversitylibrary.org/item/10277].
- Lowe, C. J., D. N. Clarke, D. M. Medeiros, D. Rokhsar, & John Gerhart. 2015. The deuterostome context of chordate origins. *Nature* 520:456–465.
- Lowe, C. J., Kuni Tagawa, Tom Humphreys, Marc Kirschner, & John Gerhart. 2004. Chapter 8. Hemichordate embryos: Procurement, culture, and basic methods. *Methods in Cell Biology* 74:171–194.
- Lowe C. J., Mark Terasaki, Michael Wu, R. M. Freeman Jr., Linda Runft, Kristen Kwan, Saori Haigo, Jochanan Aronowicz, Eric Lander, Chris Gruber, Mark Smith, Marc Kirschner, & John Gerhart. 2006. Dorsoventral patterning in hemichordates: Insights into early chordate evolution. *PLoS Biology* 13(12): e1002354. [doi.org/10.1371/journal.pbio.1002354].
- Lowe, C. J., Mike Wu, Adrian Salic, Louise Evans, Eric Lander, Nicole Stange-Thomann, C. E. Gruber, John Gerhart, & Marc Kirschner. 2003. Anteroposterior patterning in hemichordates and the origin of the chordate nervous system. *Cell* 113:853–865.
- Maletz, Jörg. 2014. Fossil Hemichordata (Pterobranchia, Enteropneusta). *Palaeogeography, Palaeoclimatology, Palaeoecology* 398:16–27

- Metschnikoff, E. 1869. Ueber Tornaria. Nachrichten von der Königlichen Gesellschaft der Wissenschaften und der Georg-Augusts Universität Göttingen 15: 287–292.
- McCoy, V. E., E. E. Saupe, J. C. Lamsdell, L. G. Tarhan, Sean McMahon, Scott Lidgard, Paul Mayer, C. D. Whalen, Carmen Soriano, Lydia Finney, Stefan Vogt, E. G. Clark, R. P. Anderson, Holger Peterman, E. R. Locatelli, & D. E. G. Briggs. 2016. The ‘Tully monster’ is a vertebrate. *Nature* 352:496–499.
- Miyamoto, Norio, & Yasunori Saito. 2007. Morphology and development of a new species of *Balanoglossus* (Hemichordata: Enteropneusta: Ptychoderidae) from Shimoda, Japan. *Zoological Science* 24:1278–1285.
- Miyamoto Norio, & Yasunori Saito. 2010. Morphological characterization of the asexual reproduction in the acorn worm *Balanoglossus simodensis*. *Development, Growth & Differentiation* 52:615–627.
- Morgan, T. H. 1891. The growth and metamorphosis of Tornaria. *Journal of Morphology* 5:407–458.
- Nanglu, Karma, J.-B. Caron, & C. B. Cameron. 2015. Using experimental decay of modern forms to reconstruct the early evolution and morphology of fossil enteropneusts. *Paleobiology* 41(3):460–478.
- Nanglu, Karma, J.-B. Caron, Simon Conway Morris, & C. B. Cameron. 2016. Cambrian suspension-feeding tubicolous enteropneusts. *BMC Biology* 14:56.
- Osborn, K. J., A. V. Gebruk, Antonina Rogacheva, & N. D. Holland. 2013. An externally brooding acorn worm (Hemichordata, Enteropneusta, Torquaratoridae) from the Russian Arctic. *The Biological Bulletin* 225(2):113–123.
- Osborn, K. J., L. A. Kuhn, I. G. Priede, Makoto Urata, A. V. Gebruk, & N. D. Holland. 2012. Diversification of acorn worms (Hemichordata, Enteropneusta) revealed in the deep sea. *Proceedings of the Royal Society (series B)* 279(1733):1646–1654.
- Packard, A. 1968. Asexual reproduction in *Balanoglossus* (Stomochordata). *Proceedings of the Royal Society, London (series B)* 171:261–272.
- Pardos, Fernando, & J. Benito. 1990. The main trunk vessels and blood components of *Glossobalanus minutus*. *European Archives of Biology (Bruxelles)* 101:455–468.
- Petersen, J. A., & A. S. F. Ditadi. 1971. Asexual reproduction in *Glossobalanus crozieri* (Ptychoderidae, Enteropneusta, Hemichordata). *Marine Biology* 9:78–85.
- Peterson, K. J., & D. J. Eernisse. 2016. The phylogeny, evolutionary developmental biology, and paleobiology of the Deuterostomia: 25 years of new techniques, new discoveries, and new ideas. *Organisms Diversity & Evolution* 16(2):401–418.
- Peterson, K. J., Y. H. Su, M. I. Arnone, B. J. Swalla, & B. L. King. 2013. MicroRNAs support the monophyly of enteropneust hemichordates. *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution* 320(6):368–374.
- Pickens, P. E. 1970. Conduction along the ventral nerve cord of a hemichordate worm. *Journal of Experimental Biology* 53:515–528.
- Priede, I. G., K. J. Osborn, A. V. Gebruk, Dan Jones, David Shale, Antonina Rogacheva, & N. D. Holland. 2012. Observations on torquaratorid acorn worms (Hemichordata, Enteropneusta) from the North Atlantic with descriptions of a new genus and three new species. *Invertebrate Biology* 131(3): 244–257.
- Punnett, R. C. 1906. The Enteropneusta. *In* J. S. Gardner, ed., *The fauna and Geography of the Maldives and Laccadive Archipelagoes*. Being the account of the work carried on and of the collections made by an expedition during the years 1899 and 1900. vol. II and supplements I and II. pl. xxvi-c, text illustrations 120-154, and index i and ii. Cambridge. Cambridge University Press. p. 631–680 + pl. xxxvii–xlvii.
- Rao, K. P. 1954. Bionomics of *Ptychodera flava* Eschscholtz (Enteropneusta). *Journal of the Madras University, B*. 24:1–5.
- Rao, K. P. 1955. Morphogenesis during regeneration in an enteropneust. *Journal of Animal Morphology and Physiology* 1:1–7.
- Ritter, W. E. 1900. Papers from the Harriman Alaska Expedition. II. *Harrimania maculosa*, a new genus and species of Enteropneusta from Alaska, with special regard to the character of its notochord. *Proceedings of the Washington Academy of Sciences* 2:111–132.
- Ritter, W. E. 1902. The movements of the Enteropneusta and the mechanism by which they are accomplished. *Biological Bulletin* 3:255–261.
- Romer, A. S. 1967. Major steps in vertebrate evolution. *Science* 158:1629–1637.
- Ruppert, E. E. 2005. Key characters uniting hemichordates and chordates: Homologies or homoplasies? *Canadian Journal of Zoology* 83:8–23.
- Ruppert, E. E., & E. J. Balse. 1986. Nephridia in the larvae of hemichordates and echinoderms. *Biological Bulletin* 171:188–196.
- Ruppert, E. E., C. B. Cameron, & J. E. Frick. 1999. Endostyle-like features of the dorsal epibranchial ridge of an enteropneust and the hypothesis of dorsal-ventral axis inversion in chordates. *Invertebrate Biology* 118(2):202–212.
- Rychel, A. L., & B. J. Swalla. 2009. Regeneration in hemichordates and echinoderms. *In* Baruch Rinkevich & Valeria Matranga, eds., *Stem Cells in Marine Organisms*. Springer. The Netherlands. p. 245–265.
- Sallan, Lauren, Sam Giles, R. S. Sansom, J. T. Clarke, Zerina Johanson, I. J. Sansom, & Philippe Janvier. 2017. The ‘Tully Monster’ is not a vertebrate: Characters, convergence and taphonomy in Palaeozoic problematic animals. *Palaeontology* 60(2):149–157.
- Satoh, Noriyuki. 2016. Chordate Origins and Evolution: The Molecular Evolutionary Road to Vertebrates. Academic Press, Elsevier. Cambridge, Massachusetts. 220 p.
- Satoh, Noriyuki, Kunifumi Tagawa, C. J. Lowe, Jr-Kai Yu, Takeshi Kawashima, Hiroki Takahashi, Michio Ogasawara, Marc Kirschner, Kanako Hisata, Yi-Hsien Su, & John Gerhart. 2014. On a possible evolutionary link of the stomochord of hemichordates to pharyngeal organs of chordates. *Genesis* 52(12):925–934.
- Schimkewitsch, W. M. 1892. On the relations between the Enteropneusta and Acrania (Russian & French).

- Publisher of the St. Petersburg Naturalist Society. 3–4:1–16.
- Seilacher, Adolf. 2007. Trace Fossil Analysis. Springer-Verlag, Berlin, Heidelberg, New York. 226 p.
- Simakov, Oleg, Takeshi Kawashima, Ferdinand Marletaz, Jerry Jenkins, Ryo Koyanagi, Therese Mitros, Kanako Hisata, Jessen Bredeson, Eiichi Shoguchi, Fuki Gyoja, Jia-Xing Yue, Yi-Chih Chen, Robert M. Freeman Jr, Akane Sasaki, Tomoe Hikosaka-Katayama, Atsuko Sato, Manabu Fujie, Kenneth W. Baughman, Judith Levine, Paul Gonzalez, Christopher Cameron, Jens H. Fritzenwanker, Ariel M. Pani, Hiroki Goto, Miyuki Kanda, Nana Arakaki, Shinichi Yamasaki, Jiaxin Qu, Andrew Cree, Yan Ding, Huyen H. Dinh, Shannon Dugan, Michael Holder, Shalini N. Jhangiani, Christie L. Kovar, Sandra L. Lee, Lora R. Lewis, Donna Morton, Lynne V. Nazareth, Geoffrey Okwuonu, Jireh Santibanez, Rui Chen, Stephen Richards, Donna M. Muzny, Andrew Gillis, Leonid Peshkin, Michael Wu, Tom Humphreys, Yi-Hsien Su, Nicholas H. Putnam, Jeremy Schmutz, Asao Fujiyama, Jr-Kai Yu, Kunifumi Tagawa, Kim C. Worley, Richard A. Gibbs, Marc W. Kirschner, Christopher J. Lowe, Noriyuki Satoh, Daniel S. Rokhsar, & John Gerhart. 2015. Hemichordate genomes and deuterostome origins. *Nature* 527:459–465.
- Smith, K. L., N. D. Holland, & H. A. Ruhl. 2005. Enteropneust production of spiral fecal trails on the deep-sea floor observed with time-lapse photography. *Deep Sea Research I* 52:1228–1240.
- Spengel, J. W. 1893. Monographie 18. Enteropneusten. Fauna und Flora des Golfes von Neapel und der angrenzenden Meeres-Abschnitte. Herausgegeben von der Zoologischen Station zu Neapel. Verlag von R. Friedländer & Sohn. Berlin. 758 p. + 37 pl. (in 2 volumes).
- Spengel, J. W. 1901. Die Benennung der Enteropneusten-Gattungen. *Zoologische Jahrbücher. Abtheilung für Systematik, Geographie und Biologie der Thiere* 15:209–218.
- Spengel, J. W. 1903. Neue Beiträge zur Kenntniss der Enteropneusten. I. *Ptychodera flava* Eschsch. von Laysan. *Zoologische Jahrbücher. Abtheilung für Anatomie und Ontogenie der Thiere* 18:271–326, pl. 24–29.
- Spengel, J. W. 1909. Pelagisches Vorkommen von Enteropneusten. *Zoologischer Anzeiger* 34:54–59.
- Stach, Thomas, & Sabrina Kaul. 2011. The postanal tail of the enteropneust *Saccoglossus kowalevskii* is a ciliary creeping organ without distinct similarities to the chordate tail. *Acta Zoologica* 92(2): 150–160.
- Strathmann, Richard, & D. Bonar. 1976. Ciliary feeding of tornaria larvae of *Ptychodera flava* (Hemichordata: Enteropneusta). *Marine Biology* 34(4):317–324.
- Szaniawski, Hubert. 2005. Cambrian chaetognaths recognized in Burgess Shale fossils. *Acta Palaeontologica Polonica* 50:1–8.
- Tagawa, Kuni. 2016. Hemichordate models. *Current Opinion in Genetics & Development* 39:71–78.
- Tagawa, Kuni, Atsuo Nishino, Tom Humphreys, & Noriyuki Satoh. 1998. The spawning and early development of the Hawaiian acorn worm (Hemichordata), *Ptychodera flava*. *Zoological Science* 15(1):85–91.
- Tassia, M. G., N. V. Whelan, & K. M. Halanych. 2017. Toll-like receptor pathway evolution in deuterostomes. *Proceedings of the National Academy of Sciences* 114(27):7055–7060.
- Tattersall, W. M. 1905. Enteropneusta from west coast of Ireland. *Annual Report of Fisheries, Ireland Scientific Investigations* 1902:213–214.
- Telford, M. J., C. J. Lowe, C. B. Cameron, Olga Ortega-Martinez, Jochanan Aronowicz, Paola Oliveri, & R. R. Copley. 2014. Phylogenomic analysis of echinoderm class relationships supports Asterozoa. *Proceedings of the Royal Society B* 281:20140479 [doi:10.1098/rspb.2014.0479].
- Thomas, I. M. 1972. Action of the gut in *Saccoglossus otagoensis* (Hemichordata: Enteropneusta). *New Zealand Journal of Marine and Freshwater Research* 6(4):560–569.
- Twedell, K. S. 1961. Regeneration of the enteropneust *Saccoglossus kowalevskii*. *Biological Bulletin* 120:118–127.
- Twitchett, R. J. 1996. The resting trace of an acorn worm (class Enteropneusta) from the Lower Triassic. *Journal of Paleontology* 70(1):128–131.
- Urata, Makoto, Sadaharu Iwasaki, & Susumu Ohtsuka. 2012. Biology of the swimming acorn worm *Glandiceps hacksii* from the Seto Inland Sea of Japan. *Zoological Science* 29(5):305–310.
- Urata, Makoto, & Masaaki Yamaguchi. 2004. The development of the enteropneust hemichordate *Balanoglossus misakiensis* Kuwano. *Zoological Science* 21(5):533–540.
- Uribe, M. E., & A. P. Larrain. 1992. Estudios biológicos en el enteropneusto *Ptychodera flava* Eschscholtz, 1825, de Bahía Concepción, Chile. I: Aspectos morfológicos y ecológicos. *Gayana. Zoología* 56:141–180.
- Wagner, Nicolas. 1885. Die Wirbellosen des Weissen Meeres. *Zoologische Forschungen an der Küste des Solowetzischen Meerbusens in den Sommermonaten der Jahre 1877, 1878, 1879 und 1882. Erster Band mit 22 zum Theil farbigen Tafeln und mehreren Holzschnitten.* Verlag Wilhelm Engelmann. Leipzig. 1-171, 21 pl.
- Walcott, C. D. 1911. Cambrian Geology and Paleontology II. Middle Cambrian annelids. *Smithsonian Miscellaneous Collections* 57(5):109–445.
- Walcott, C. D. 1931. Addenda to descriptions of Burgess Shale fossils. *Smithsonian Miscellaneous Collections* 85(3):1–46.
- Welsch, Ulrich, & Volker Storch. 1970. The fine structure of the stomochord of the enteropneusts *Harrimania kupfferi* and *Ptychodera flava*. *Zeitschrift für Zellforschung* 107:234–239.
- Willey, Arthur. 1898. *Spengelia*, a new genus of Enteropneusta. *Quarterly Journal of Microscopical Science* 40:623–630.
- Willey, Arthur. 1899. Remarks on some recent work on the Protochordata, with a condensed account of some fresh observations on the Enteropneusta. *Quarterly Journal of Microscopical Science* 42:223–244.

- Winchell, C. J., Jack Sullivan, C. B. Cameron, B. J. Swalla, & Jon Mallatt. 2002. Evaluating hypotheses of deuterostome phylogeny and chordate evolution with new LSU and SSU ribosomal DNA data. *Molecular Biology and Evolution* 19(5):762–776.
- Woodin, S. A., M. D. Walla, & D. E. Lincoln. 1987. Occurrence of brominated compounds in soft-bottom benthic organisms. *Journal of Experimental Marine Biology and Ecology* 107:209–217.
- Woodwick, K. H., & Terry Sensenbaugh. 1985. *Saxipendium coronatum*, new genus, new species (Hemichordata: Enteropneusta): The unusual spaghetti worms of the Galapagos Rift hydrothermal vents. *Proceedings of the Biological Society of Washington* 98(2):351–365.
- Worsaae, Katrine, Wolfgang Sterrer, Sabrina Kaul-Strehlow, Anders Hay-Schmidt, & Gonzalo Giribet. 2012. An anatomical description of a miniaturized acorn worm (Hemichordata, Enteropneusta) with asexual reproduction by paratomy. *Plos One* 7(11):1–19.
- Yoshimatu, S., & Teruaki Nishikawa. 1999. Swimming swarms of a usually benthic enteropneust. *Glandiceps* sp. in the Seto Inland Sea, Japan, found in 1998. *Proceedings of the Seventieth Annual Meeting of the Zoological Society of Japan. Zoological Science* 16:39.